

Nonresponse to Hepatitis B Vaccines and the Kinetics of Anti-HBs Production

Jane N. Zuckerman

Academic Unit of Travel Medicine and Vaccines, The Royal Free Hospital School of Medicine, London, United Kingdom

No empirical data are available for the anti-HBs titre required for protection against infection with the hepatitis B virus (HBV), but nonresponders to hepatitis B vaccines remain susceptible to infection. There may be an unexplained qualitative difference between hyporesponders and true nonresponders, but there is a clear association with HLA haplotypes. There is no evidence for silent infection with HBV in nonresponders.

© 1996 Wiley-Liss, Inc.

KEY WORDS: Hepatitis B, vaccines, nonresponse

INTRODUCTION

Systematic vaccination of individuals at risk of exposure to hepatitis B virus (HBV) has been the main method of controlling the morbidity and mortality associated with hepatitis B infection. The first hepatitis B vaccine was manufactured by the purification and inactivation of hepatitis B surface antigen (HBsAg) obtained from the plasma of chronic HBV carriers [Szmuness et al., 1981; Hadler et al., 1986; Jilg et al., 1988]. This was soon followed by the production of HBsAg using recombinant DNA techniques and expression of the S antigen component in yeast cells.

All studies of the antibody response to currently licensed plasma-derived hepatitis B vaccines and hepatitis B vaccines prepared by recombinant DNA technology have shown that between 5% and 10% or more of healthy immunocompetent subjects do not mount an antibody response (anti-HBs) to the surface antigen component (HBsAg) present in these preparations (nonresponders) or that they respond poorly (hyporesponders) [Dienstag et al., 1984; Craven et al., 1986; Westmoreland et al., 1990; Wood et al., 1993]. The exact proportion depends partly on the definition of nonresponsiveness or hyporesponsiveness, generally less than 10 IU/L or 100 IU/L of anti-HBs, respectively, against an international antibody standard.

Nonresponders remain susceptible to infection with HBV. While several factors are known to affect adversely the antibody response to HBsAg including the

site and route of injection, gender, advancing age, body mass (overweight), immunosuppression and immunodeficiency, and mechanisms underlying nonresponsiveness to the S component of HBsAg in humans remain largely unexplained, evidence is accumulating that there is an association between different HLA-DR alleles and specific low responsiveness in different ethnic populations. Considerable experimental evidence is available to suggest that the ability to produce antibody in response to specific protein antigens is controlled by dominant autosomal class II genes of the major histocompatibility complex (MHC) in the murine model [reviewed in Alper et al., 1989; Milich, 1991; Kruskall et al., 1992]. Much effort has been devoted to overcoming class II-linked nonresponsiveness to current hepatitis B vaccine [e.g., Milich et al., 1985a, 1986; Arif et al., 1988].

There is evidence that the pre-S1 and pre-S2 domains have an important immunogenic role in augmenting anti-HBs responses, preventing the attachment of the virus to hepatocytes and eliciting antibodies which are effective in viral clearance, stimulating cellular immune responses, and circumventing genetic nonresponsiveness to the S antigen [Milich et al., 1985a; Klinkert et al., 1986; Alberti et al., 1988; Gerlich et al., 1990]. Thus, a number of studies indicated that the inclusion of pre-S components in recombinant or future synthetic vaccines should be developed. For example, the pre-S2 region is more immunogenic at the T and B-cell levels than the S regions in the mouse model [Milich et al., 1985a,b], as is the case with pre-S1 in the mouse [Milich et al., 1986] and in humans [Ferrari et al. 1992], and circumvents S region nonresponsiveness at the level of antibody production.

Indeed, Milich et al. [1986] demonstrated in the murine model that the independence of MHC-linked gene regulation of immune responses to pre-S1, pre-S2, and S regions of HBsAg would assure fewer genetic nonresponders to a vaccine containing all three antigenic

Accepted for publication July 17, 1996.

Address reprint requests to Dr. Jane N. Zuckerman, Academic Unit of Travel Medicine and Vaccines, Royal Free Hospital School of Medicine, Rowland Hill Street, London NW3 2PF, UK.

regions. Studies conducted in humans with experimental recombinant hepatitis B vaccines containing all three S components of the viral envelope polypeptides demonstrated the enhanced immunogenicity of such preparations when compared with conventional yeast-derived vaccines [Yap et al., 1992, 1995; Shouval et al., 1994; Zuckerman et al., 1996], although several earlier studies with vaccines containing the S and pre-S components revealed significant differences from preparations containing only the S antigen [Marescot et al., 1989; Ferrari et al., 1992; Clements et al., 1994; Suzuki et al., 1994].

No empirical data are available for the anti-HBs titre required for protection against particular routes of infection or the size of the infectious inoculum. The minimum protective level following immunisation has been set in earlier protective efficacy studies at 10 IU/L or more of anti-HBs [Szmuness et al., 1981; Francis et al., 1982]. In both studies most cases of HBV infection occurred in subjects who mounted little or no anti-HBs response. Specifically, a protective level of anti-HBs was defined as 10 IU/L against an international standard [Stevens et al., 1984a; Centers for Disease Control, 1987]. Various studies have also demonstrated that the risk of HBV infection increases as anti-HBs levels decline to 10 IU/L [Stevens et al., 1984b; Coursaget et al., 1986; Hadler et al., 1986; Taylor and Stevens, 1988]. For example, Hadler et al. [1986] reported in a follow-up study of vaccinated homosexual men an overall incidence of HBV infection of 2.9 per 100 person-years with nearly 75% occurring in subjects with anti-HBs titres <10 IU/L at the time of infection and in only a few with anti-HBs titres >50 IU/L. A lower and asymptomatic infection rate of 0.8 per 100 person-years was observed after immunisation of health care workers in nephrology units who had antibody titres of <50 IU/L [Couroucé et al., 1988].

THE KINETICS OF ANTI-HBs RESPONSE

The titre of vaccine-induced anti-HBs declines, often rapidly, during the months and years following immunisation. The highest anti-HBs titres are generally observed 1 month after booster vaccination followed by rapid decline during the next 12 months and thereafter more slowly [see for example Hilleman, 1984; Jilg et al., 1984; Ambrosch et al., 1987; Gibas et al., 1988; Nommensen et al., 1989; Wiseman et al., 1989]. Mathematical models were designed and an equation was derived consisting of several exponential terms with different half-life periods. It is considered by some researchers that the decline of anti-HBs concentration in an immunised subject can be predicted accurately by such antibody kinetics and preliminary recommendations before the next booster vaccination [Jilg et al., 1984; Fagan et al., 1984a,b; Ambrosch et al., 1987; Nommensen et al., 1989]. If the minimum protective level is accepted at 10 IU/L, which is being debated, consideration should be given to the diversity of the individual immune response and the decrease in levels of anti-HBs as well as to possible errors in quantitative anti-

HBs determinations. It would then be reasonable to define a level of >10 IU/L and <100 IU/L as an indication for booster immunisation. It has been demonstrated that a booster inoculation results in a rapid increase in anti-HBs titres within 4 days [Jilg et al., 1988]. However, even this time delay might permit infection of hepatocytes [Nommensen et al., 1989].

Several options are therefore under consideration for maintaining protective immunity against hepatitis B infection:

1. Relying upon immunological memory to protect against clinical infection and its complications [Centers for Disease Control, 1991; Tilzey, 1995], a view which is supported by *in vitro* studies showing immunological memory for HBsAg in B cells derived from vaccinated subjects who have lost their anti-HBs, but not in B cells from nonresponders [van Hattum et al., 1991], and indeed one cannot recall what has never been memorised [McIntyre, 1995].

2. Providing booster vaccination to all vaccinated subjects at regular intervals without determination of anti-HBs. This option is not supported by a number of investigators because nonresponders must be detected [McIntyre, 1995; Tedder et al., 1993]. Also, while an anti-HBs titre of about 10 IU/L may in theory be protective, this level is not protective from a laboratory point of view since many serum samples may give nonspecific reactions at this antibody level [Westmoreland et al., 1990; Tedder et al., 1993].

3. Testing anti-HBs levels 1 month after the first booster and administering the next booster before the minimum protective level is reached, which is the preferred option. A protective level of 100 IU/L seems to be appropriate.

REACTIVATION IN NONRESPONDERS

There are numerous reports in the literature that the administration of four, five, or six or more doses of vaccine in apparent nonresponders or hyporesponders resulted in the production of anti-HBs in as many as 50% [Craven et al., 1986; Fagan et al., 1987b; Pasko et al., 1990; Westmoreland et al., 1990], although most reports concern a small number of subjects. A study of some 26 hyporesponders and nonresponders in The Netherlands [Wisnans et al., 1988] revealed that some developed anti-HBs after up to a further six inoculations while others failed to respond, leading to a comment that there is an (unexplained) qualitative difference between hyporesponders and "real" nonresponders. Several other explanations had been offered, apart from the administration of the vaccine into an inappropriate site such as the gluteal muscles or intradermally or improper storage of the vaccine. A "slow response" has been considered, as well as modulation of the immune response by genetic factors particularly in relation to HLA type, nonspecific cellular immune defects, concurrent infections, and preferential stimulation of T-suppressor cells.

HLA HAPLOTYPES IN NONRESPONDERS TO HEPATITIS B VACCINE AND IN RESPONSE TO (HEP B-3) A NEW S, PRE-S1, AND PRE-S2 VACCINE

The HLA class I and class II alleles were determined by McDermott et al. [1996] in the 86 vaccine nonresponders participating in a study of the Hep B-3 vaccine [Zuckerman et al., 1996] by a lymphomicrocytotoxicity technique for class I HLA typing using commercially available plates (Biotest, Germany), and by polymerase chain reaction (PCR) for class II HLA typing. A control group of 115 subjects who were hepatitis B vaccine responders and an additional group of 125 Caucasian new volunteer donors at the Anthony Nolan Bone Marrow Trust were included as further control subjects.

A significant association was found between the HLA phenotype B44;DRB1*0701; DQB1*0201 and the vaccine nonresponse ($P = 0.02$). Those with the phenotype B44; DRB1*0701; DQB1*0201 are nearly four times more likely to be antibody nonresponders when compared to hepatitis B vaccine responders ($P = 0.01$). Previous studies have identified the association of the extended haplotypes B44; DR7; FC31; and B8; DR3; SC01 based on six of nine individuals who were nonresponders to the S vaccine. The correlation of B44; DRB1*0701; DQB1*0201 with hepatitis B vaccine nonresponse was confirmed in the present study representing more accurately the frequency of HLA in the Caucasian population. Despite this, we could not identify a significant association with the HLA phenotype B8; DR3 with non-response to HBsAg vaccination as described previously. The frequency of the HLA allele DQB1*0201 was high in our nonresponder population ($P = 0.002$). This allele is in strong linkage disequilibrium with HLA alleles DRB1*0701 and DRB1*0301 which are found in 85% of vaccine nonresponders. The HLA alleles DRB1*1501 and DRB1*0101/02/03 were found at a lower frequency in the nonresponders when compared to the HLA and vaccine control groups, respectively, but were not significant when adjusted for the number of tests carried out. The DR1 antigen has been described previously by less stringent analysis [Walker et al., 1981]. Interestingly, the HLA allele DQB1*0602 was represented at a significantly low frequency in the nonresponder population ($P = 0.02$). This allele is in linkage with DRB1*1501, suggesting that there was an association between the molecular subtypes of DR1 and the response to hepatitis B vaccination.

In summary, a high frequency of HLA class II allele DRB1*0701 and the phenotype B44; DRB1*0701; DQB1*0201 was found in nonresponders compared to controls [McDermott et al., 1996]. There were low frequencies of DRB1*1501, DQB1*0602, and phenotype DRB1*1501; DQB1*0602 among nonresponders. All the initial nonresponders expressing the phenotype B7; DRB1*1501; DQB1*0602 responded to the new vaccine with antibody titres >100 IU/L. The majority of those who failed to mount an antibody response expressed

the phenotypes B8; DRB1*0301; DQB1*0201 or B44; DRB1*0701; DQB1*0201 [McDermott et al., 1996].

Nonresponse to the additional pre-S1 and pre-S2 components in the vaccine may be due to dysfunctional interaction of specific HLA molecules with these antigens, lack of T-cell activation, or deletion of specific T cells during thymic education. Further study and typing for additional polymorphism within the HLA-mediated immune response may provide an insight into the mechanism of failure to respond to immunisation against hepatitis B [McDermott et al., 1996]. However, it should be noted that immunogenetic analysis confirmed that an initially distinct group of nonresponders was indeed included in the study of the new pre-S1, pre-S2, S vaccine.

NONRESPONDERS AND SUSCEPTIBILITY TO INFECTION WITH HBV

Whilst it is accepted that about 5–15% of fully immunocompetent healthy individuals do not mount a humoral antibody response to currently available hepatitis B vaccines, and others are poor responders, there is little in the more recent literature based on long-term follow-up to address the issue of whether such persons are susceptible to infection with HBV. The kinetics of antibody and the issue of post-vaccination testing have been discussed above, and it is undoubtedly of importance for health care workers, their patients, and their employers (under current UK Department of Health Guidelines) to be aware of their protection or lack of protection after immunisation. Equal reliance on an adequate protective cell-mediated immune response (which in the vast majority has not been measured) or primed (or otherwise) immunological memory to mount an anamnestic response may not be entirely satisfactory.

Further, it is easy to be too dismissive of the importance of symptomless HBV infection based on anti-HBc seroconversion in vaccine responders and nonresponders. The suggestion in an editorial [Hall, 1993] that “whether antibody responses after vaccination should be verified and subsequent decay documented will depend on local resources” is not acceptable in the interest of the public health, apart from other considerations.

An early placebo-controlled study was carried out with a plasma-derived vaccine in an HBV “high-risk” setting in 353 staff, patients on maintenance haemodialysis, and their relatives in France in 1975 [Maupas et al., 1979]. Follow-up of 73 patients and 191 staff showed that vaccinated subjects who did not respond to the vaccine by developing anti-HBs were infected at the same rate as the unvaccinated controls, i.e., nearly 50% as indicated either by anti-HBc production alone (5%), transient antigenaemia (15%), or prolonged antigenaemia (25%). Many of the subjects who developed infection within 2 months of immunisation were patients, who tend to mount a delayed or slow anti-HBs response and were likely to be incubating the infection. Thirteen staff members (60%) were nonresponders and

nine became infected with HBV within 4 to 12 months after the first inoculation.

It should be noted that interpretation of parts of the report is difficult. Other studies [Stevens et al., 1984a; Coursaget et al., 1986; Hadler et al., 1986; Couroucé et al., 1988; Taylor and Stevens, 1988] have shown that the risk of HBV infection increases as anti-HBs levels decline to 10 IU/L in responders. There are few reports concerning nonresponders. Nevertheless, the initial efficacy trials of the plasma-derived hepatitis B vaccine (produced by Merck, Sharp & Dohme in the United States) provide evidence of the continuing susceptibility of persons who receive a complete course of vaccine but develop less than 10 IU/L of anti-HBs. For example, the study conducted by Szmuness et al. [1981] revealed that 7 of 21 (33%) of vaccinated nonresponder male homosexuals became infected during an 18-month period of surveillance. That compared with 92 of 426 (22%) placebo recipients infected during the same period. The evaluation in another study of long-term protection by hepatitis B vaccine for 7–9 years revealed 36 HBV infections among 139 male homosexuals who had no detectable anti-HBs after three doses of vaccine [Hadler et al., 1991]. In an earlier trial, the same investigators noted that HBV infection occurred in 55 vaccinated subjects with a poor antibody response and 2 became carriers of HBV, both of whom were nonresponders [Hadler et al., 1986]. In another study there were 4 “vaccine failures” among 15 babies born to high-risk mothers; one infant nonresponder became infected at the age of 10 months and one poor responder became infected at the age of 6.5 months and remained *e* antigen positive for 5 months of the follow-up [Flower and Tanner, 1988].

There are apparently no reports of a cohort of healthy nonresponders to vaccination who have been surveyed systematically for a sufficient number of person-years to closely estimate susceptibility to infection. It is proposed to follow-up by serological surveillance the 86 nonresponder participants in the Hep B-3 vaccine, amongst whom the overall rate of seroconversion in terms of anti-HBs titre of >10 IU/L was 66%, with rates ranging from 55 to 76% across antigen doses of 5, 10, 20, and 40 µg [Zuckerman et al., 1996].

NONRESPONDERS AND SILENT INFECTION

A brief report [Lou et al., 1992] noted that 6.4% of 214 subjects in China who were immunised with the Merck, Sharp & Dohme hepatitis B vaccine and 12.5% of 96 subjects who received a locally produced vaccine did not respond. HBV DNA was detected by PCR in over 60% of the nonresponders in each group, suggesting that nonresponsiveness to hepatitis B vaccine may be due to immunotolerance or immunosuppression induced by latent HBV infection. Other reports suggested that HBV *e* antigen can cause immunotolerance and chronic HBV [Brunetto et al., 1991], and that HBV itself may cause immunotolerance by infecting directly T and B lymphocytes resulting in viral persistence [Oldstone, 1989], or through different mechanisms triggered

by viral infection leading to imbalance in immunoregulation [Paller and Mallory, 1991].

No evidence of latent HBV infection was found in 86 nonresponders in the study carried out by Zuckerman et al. [1996]. This observation was made after the repeated absence of serological markers of infection in the subjects (anti-HBc and HBsAg), and by the absence of HBV DNA by nested-PCR in many of the nonresponder subjects at the Royal Free Hospital and School of Medicine (Dr. T.J. Harrison, personal communication) and in four volunteers with anti-HBc.

ATTEMPTS TO OVERCOME NONRESPONSIVENESS BY THE USE OF IMMUNOMODULATORS

Attempts have been made to enhance the anti-HBs response following immunisation, particularly in patients treated by maintenance haemodialysis, but often with conflicting results or in limited studies, which have not been confirmed: alpha interferon [Grob et al., 1984; Goldwater, 1994]; interleukin-2 [Meuer et al., 1989; Jungers et al., 1994]; and thymopentin [Zaruba et al., 1983; Melappioni et al., 1992]. Other substances such as experimental oral adjuvants in mice and oestrogen have been referred to for the sake of completion.

ACKNOWLEDGMENTS

The Hep B-3 vaccine study in nonresponders was carried out in collaboration with the WHO Collaborating Centre for Reference and Research on Viral Diseases at the Royal Free Hospital School of Medicine, and with Dr. A. Williams and Dr. Fiona Craig, Medeva Plc. The study of HLA haplotypes is part of a continuing collaborative study with Dr. A. Madrigal and Mr. A.B. McDermott of the Anthony Nolan Research Institute of the Anthony Nolan Bone Marrow Trust, Royal Free Hospital, London.

REFERENCES

- Alberti A, Cavalletto D, Pontisso P, Chemello L, Tagariello G, Belussi F (1988): Antibody response to pre-S2 and hepatitis B virus induced liver damage. *Lancet* I:1421–1424.
- Alper CA, Kruskall MS, Marcus-Bagley D, Craven DE, Katz AJ, Brink SJ, Dienstag JL, Awdeh Z, Yunis EJ (1989): Genetic prediction of nonresponse to hepatitis B vaccine. *The New England Journal of Medicine* 321:708–712.
- Ambrosch F, Frisch-Niggemeyer W, Kremsner P, Kunz Ch, Andre F, Safary A, Wiedermann G (1987): Persistence of vaccine-induced antibodies to hepatitis B surface antigen and the need for booster vaccination in adult subjects. *Postgraduate Medical Journal* 63(Suppl 2):129–135.
- Arif M, Mitchison NA, Zuckerman AJ (1988): Genetics of nonresponders to hepatitis B surface antigen and possible ways of circumventing “nonresponse.” In Zuckerman AJ (ed.): “Viral Hepatitis and Liver Disease.” New York: Alan R. Liss, pp 714–716.
- Brunetto MR, Giardin MM, Oliveri F, Chiaberge E, Baldi M, Alfarano A, Serra A, Saracco G, Verme G, Will H, Bonino F (1991): Wild-type and *e* antigen-minus hepatitis B viruses and course of chronic hepatitis. *Proceedings of the National Academy of Sciences, USA* 88:4186–4190.
- Centers for Disease Control (1991): Hepatitis B virus: A comprehensive strategy for eliminating transmission in the United States through universal childhood vaccination. *Morbidity & Mortality Weekly Report* 40(RR-13):1–19.
- Clements ML, Miskovsky E, Davidson M, Cupps T, Kumwenda N, Sandman LA, West D, Hesley T, Loli V, Miller W, Calandra G,

- Krugman S (1994): Effect of age on the immunogenicity of yeast recombinant hepatitis B vaccines containing surface antigen (S) or pre-S2 + S antigens. *Journal of Infectious Diseases* 170:510-516.
- Couroucé A-M, Laplanche A, Benhamou E, Jungers P (1988): Long-term efficacy of hepatitis B vaccine in healthy adults. In Zuckerman AJ (ed.): "Viral Hepatitis and Liver Disease." New York: Alan R. Liss, pp 1002-1005.
- Coursaget P, Yvonnet B, Chotard J-N, Sarr M, Vincelot P, N'Doye R, Diop-Mar I, Chiron JP (1986): Seven-year study of hepatitis B vaccine efficacy in infants from an endemic area (Senegal). *Lancet* 2:1143-1145.
- Craven DE, Awdeh ZL, Kunches LM, Yunis EJ, Dienstag JL, Werner BG, Polk F, Snyderman DR, Platt R, Crumpacker CS, Grady GF, Alper CA (1986): Nonresponsiveness to hepatitis B vaccine in health care workers. *Annals of Internal Medicine* 105:356-360.
- Dienstag JL, Werner BG, Polk F, Snyderman DR, Craven DE, Platt R, Crumpacker CS, Ouellet-Hellstrom R, Grady GF (1984): Hepatitis B vaccine in health care personnel: Safety, immunogenicity, and indicators of efficacy. *Annals of Internal Medicine* 101:34-40.
- Fagan EA, Trolley P, Smith H, Peters MP, Coleman J, Elliott P, Williams R, Eddleston ALWF (1987a): Hepatitis B vaccine: Immunogenicity and follow-up including two year boosters doses in high risk health personnel in a London teaching hospital. *Journal of Medical Virology* 21:49-56.
- Fagan E, Williams R, Eddleston ALWF (1987b): Hepatitis B vaccine: Responder status and timing of additional booster doses. *Lancet* 2:561.
- Ferrari C, Cavalli A, Penna A, Valli A, Bertolotti A, Pedretti G, Pilli M, Vitali P, Neri TM, Giuberti T, Fiaccadori F (1992): Fine specificity of the human T-cell response to the hepatitis B virus preS1 antigen. *Gastroenterology* 103:255-263.
- Flower AJE, Tanner MS (1988): Use of hepatitis B vaccine in babies of hepatitis B surface antigen-positive carrier mothers. In Zuckerman AJ (ed): "Viral Hepatitis and Liver Disease." New York; Alan R. Liss, pp 980-981.
- Francis DP, Hadler SC, Thompson SE, Maynard JE, Ostrow DG, Altman N, Braff EH, O'Malley P, Hawkins D, Judson FN, Penley K, Nylund T, Christie G, Meyers F, Moore JN, Gardner A, Doto IL, Miller JH, Reynolds GH, Murphy BL, Shable CA, Clark BT, Curren JW, Redeker AG (1982): The prevention of hepatitis B with vaccine. *Annals of Internal Medicine* 97:362-366.
- Gerlich WH, Deepen R, Heermann KH, Krone B, Lu XY, Seifer M, Thomssen R (1990): Protective potential of hepatitis B virus antigens other than the S gene protein. *Vaccine* 8:S63-S68.
- Gibas A, Watkins E, Hinkle C, Dienstag J (1988): Long term persistence of protective antibody after hepatitis B vaccination of healthy adults. In Zuckerman AJ (ed): "Viral Hepatitis and Liver Disease." New York: Alan R. Liss, pp 998-1001.
- Goldwater PN (1994): Randomized comparative trial of interferon-alpha versus placebo in hepatitis B vaccine non-responders and hyporesponders. *Vaccine* 12:410-414.
- Grob PJ, Joller-Jemelka HI, Binswanger U, Zaruba K, Descocudres C, Fernex M (1984): Interferon as an adjuvant for hepatitis B vaccination in non and low-responder populations. *European Journal of Clinical Microbiology* 3:195-198.
- Hadler SC, Francis DP, Maynard JE, Thompson SE, Judson FN, Echenberg DF, Ostrow DG, O'Malley PM, Penley KA, Altman NL, Braff E, Shipman GF, Coleman PJ, Mandel EJ (1986): Long-term immunogenicity and efficacy of hepatitis B vaccine in homosexual men. *New England Journal of Medicine* 315:209-214.
- Hadler SC, Coleman PJ, O'Malley P, Judson FN, Altman N (1991): Evaluation of long-term protection by hepatitis B vaccine for seven to nine years in homosexual men. In Hollinger FB, Lemon SB, Margolis HS (eds). "Viral Hepatitis and Liver Disease." Baltimore: Williams & Wilkins, pp 766-768.
- Hall AJ (1993): Hepatitis B vaccination: Protection for how long and against what? *British Medical Journal*, 307:276-277.
- Hilleman MR (1984): Immunologic prevention of human hepatitis. *Perspective in Biology and Medicine*, 27:543-557.
- Jilg W, Schmidt M, Deinhardt F, Zachoval R (1984): Hepatitis B vaccination: How long does protection last? *Lancet* 2:458.
- Jilg W, Schmidt M, Deinhardt F (1988): Immune response to hepatitis B revaccination. *Journal of Medical Virology* 24:377-384.
- Jungers P, Devillier P, Salomon H, Cerisier JE, Couroucé AM (1994): Randomised placebo-controlled trial of recombinant interleukin-2 in chronic uraemic patients who are non-responders to hepatitis B vaccine. *Lancet* 344:856-857.
- Klinkert M, Theilmann L, Pfaff E, Schaller H (1986): Pre-S antigens and antibodies early in the course of acute hepatitis B virus infection. *Journal of Virology* 52:522-525.
- Kruskall MS, Alper CA, Awdeh Z, Yunis EJ, Marcus-Bagley D (1992): The immune response to hepatitis B vaccine in humans: Inheritance patterns in families. *Journal of Experimental Medicine* 175:495-502.
- Lou K-X, Wang L-P, Nie J, Jiang S (1992): Is non-responsiveness to hepatitis B vaccine due to latent hepatitis B virus infection? *Journal of Infectious Diseases* 165:777-778.
- Marescot MR, Budkowska A, Pillot J, Debre P (1989): HLA linked immune response to S and pre-S2 gene products in hepatitis B vaccination. *Tissue Antigens* 33:495-500.
- Maupas Ph, Goudeau A, Coursaget P, Drucker J, Barin F, Andre M (1979): Immunization against hepatitis B in man: A pilot study of two years' duration. In Vyas GN, Cohen SN, Schmid R (eds): "Viral Hepatitis: A Contemporary Assessment of Etiology, Epidemiology, Pathogenesis and Prevention." Philadelphia: Franklin Institute Press, pp 359-556.
- McDermott AB, Zuckerman JN, Sabin CA, Madrigal JA (1996): HLA haplotypes in non-responders to hepatitis B vaccine and in response to a novel recombinant vaccine. (Submitted).
- McIntyre PJ (1995): Hepatitis B vaccination follow-up. *Lancet* 345:1575.
- Melapponi M, Baldassari M, Saldini S, Radicioni R, Panichi N (1992): Use of immunomodulators (Thymopentin) in hepatitis B vaccine in elderly patients undergoing chronic hemodialysis. *Nephron* 61:358-359.
- Meuer SC, Dumann H, Meyer zum Buschenfelde K-H, Kohler H (1989): Low dose interleukin-2 induces systemic immune responses against Hbs Ag in immunodeficient non-responders to hepatitis B vaccination. *Lancet* 1:15-17.
- Milich DR (1991): Immune response to hepatitis B virus proteins: Relevance of the murine model. *Seminars in Liver Disease* 11:93-112.
- Milich DR, McNamara NK, McLachlan A, Thornton GB, Chisari FV (1985a): Distinct H-2 linked regulation of T-cell responses to the Pre-S and S regions of the same hepatitis B surface polypeptide allows circumvention of nonresponsiveness to the S region. *Proceedings of the National Academy of Sciences USA* 82:8168-8172.
- Milich DR, Thornton GB, Neurath AR, Kent SB, Michel M-L, Tiollais P, Chisari FV (1985b): Enhanced immunogenicity of the Pre-S region of hepatitis B surface antigen. *Science* 228:1195-1199.
- Milich DR, McLachlan A, Chisari FV, Kent SB, Thornton GB (1986): Immune response to the pre-S(1) region of hepatitis B surface antigen (HBsAg): A pre-S(1)-specific T cell response can bypass nonresponsiveness to the pre-S(2) and S regions of the HBsAg. *Journal of Immunology* 137:315-322.
- Nommensen FE, Go ST, MacLaren DM (1989): Half-life of HBs antibody after hepatitis B vaccination: An aid to timing of booster vaccination. *Lancet* 2:847-849.
- Oldstone MBA (1989): Viral persistence. *Cell* 56:517-552.
- Paller AS, Mallory SB (1991): Acquired forms of immunosuppression. *Journal of the American Academy of Dermatology* 24:482-488.
- Pasko MT, Pharm D, Beam TR Jr (1990): Persistence of anti-Hbs among health care personnel immunized with hepatitis B vaccine. *American Journal of Public Health* 80:590-593.
- Shouval D, Ilan Y, Adler R, Deepen R, Panet A, Even-Chen Z, Gorecki M, Gerlich WH (1994): Improved immunogenicity in mice of a mammalian cell-derived recombinant hepatitis B vaccine containing pre-S1 and pre-S2 antigens as compared with conventional yeast-derived vaccines. *Vaccine* 12:1453-1459.
- Stevens CE, Alter HJ, Taylor PE, Zaig EA, Harley EJ, Szmunn W (1984a): Hepatitis B vaccine in patients receiving haemodialysis: Immunogenicity and efficacy. *New England Journal of Medicine* 311:496-501.
- Stevens CE, Taylor PE, Tong MJ, Toy PT, Vyas GN (1984b): Hepatitis B vaccine: An overview. In Vyas GN, Dienstag JL, Hoofnagle JH (eds): "Viral Hepatitis and Liver Disease." Orlando: Grune and Stratton, pp 275-291.
- Suzuki H, Lino S, Shiraki K, Akahane Y, Okamoto H, Domoto K, Mishihiro S (1994): Safety and efficacy of a recombinant yeast-derived pre-S2 +S-containing hepatitis B vaccine (TGP-943): Phase I, 2, and 3 clinical testing. *Vaccine* 12:1090-1095.
- Szmunn W, Stevens CE, Zang EA, Harley EJ, Kellner A (1981): A controlled clinical trial of the efficacy of the hepatitis B vaccine (Heptavax B). A Final Report. *Hepatology* 1:377-385.

- Taylor PE, Stevens CE (1988): Persistence of antibody to hepatitis B surface antigen after vaccination with hepatitis B vaccine. In Zuckerman AJ (ed): "Viral Hepatitis and Liver Disease." New York: Alan R. Liss, pp 995-997.
- Tedder RS, Zuckerman MA, Brink N (1993): Hepatitis B vaccination: Non-responders must be detected. *Lancet* 307:732.
- Tilzey AJ (1995): Hepatitis B vaccine boosting; the debate continues. *Lancet* 345:1000-1001.
- van Hattum J, Maikoe T, Poel J, de Gast GC (1991): In vitro anti-HBs production by individual B cells of responders to hepatitis B vaccine who subsequently lost antibody. In Hollinger BF, Lemon SM, Margolis H (eds): "Viral Hepatitis and Liver Disease." Baltimore: Williams and Wilkins, pp 774-776.
- Walker M, Szmuness W, Stevens CE, Rubenstein P (1981): Genetics of anti-HBs responsiveness: HLA DR7 and nonresponsiveness to hepatitis B vaccination. *Transfusion* 21:601.
- Westmoreland D, Player V, Heap DC, Hammond A (1990): Immunization against hepatitis B-what can we expect? *Epidemiological Infection* 104:499-509.
- Wiseman P, van Hattum J, Mudde GC, Endeman HJ, Poel J, de Gast GC (1989): Is booster injection with hepatitis B vaccine necessary in healthy responders? A study of the immune response. *Journal of Hepatology* 8:236-240.
- Wiseman P, van Hattum J, Stelling T, Poel J, De Gast GC (1988): Effect of supplementary vaccination in healthy non-responders to hepatitis B vaccination. *Hepatogastroenterology* 35:78-79.
- Wood RC, MacDonald KL, White KE, Hedberg CW, Hanson M, Osterholm MT (1993): Risk factors for lack of detectable antibody response following hepatitis B vaccination of Minnesota health care workers. *Journal of the American Medical Association* 270:2935-2939.
- Yap I, Guan R, Chan SH (1995): Study on the comparative immunogenicity of a recombinant DNA hepatitis B vaccine containing pre-S components of the HBV coat protein with non pre-S containing vaccines. *Journal of Gastroenterology and Hepatology* 10:51-55.
- Yap I, Guan R, Chan SH (1992): Recombinant DNA hepatitis B vaccine containing pre S-components of the HBV coat protein—a preliminary study of immunogenicity. *Vaccine* 10:439-442.
- Zaruba K, Rastorfer M, Grob PJ, Joller-Jemelka H, Bolla K (1983): Thymopentin as adjuvant in non-responders or hyporesponders to hepatitis B vaccination. *Lancet* 2:1245.
- Zuckerman JN, Craig FN, Williams A, Zuckerman AJ (1996): Immunogenicity of a novel triple-S hepatitis B vaccine in nonresponder health care workers. Submitted for publication.